

# Lake Ontario Nearshore Water Quality Atlas

1976 - 79



Ministry  
of the  
Environment

Hon. Harry C. Parrott, D.D.S.,  
*Minister*

Graham W. S. Scott, Q.C.,  
*Deputy Minister*

### Copyright Provisions and Restrictions on Copying:

This Ontario Ministry of the Environment work is protected by Crown copyright (unless otherwise indicated), which is held by the Queen's Printer for Ontario. It may be reproduced for non-commercial purposes if credit is given and Crown copyright is acknowledged.

It may not be reproduced, in all or in part, for any commercial purpose except under a licence from the Queen's Printer for Ontario.

For information on reproducing Government of Ontario works, please contact ServiceOntario Publications at [copyright@ontario.ca](mailto:copyright@ontario.ca)

# Lake Ontario Nearshore Water Quality Atlas

1976 – 79

Toronto, Ontario, 1980



Ontario

Ministry  
of the  
Environment

Water Resources Branch  
Great Lakes Unit

© 1980 Her Majesty the Queen in Right of Ontario

Additional copies of this atlas may be obtained in person from  
the Ontario Government Bookstore, 880 Bay Street, Toronto, Ontario,  
or by mail from the Ontario Government Publications Centre,  
880 Bay Street, 5th Floor, Toronto, Ontario, M5S 1Z8

ISSN 0227-1028

Parts of this publication may be republished if credit is given.



# CONTENTS

	Page		Page
INTRODUCTION .....	1	Conductivity .....	11
GUIDE TO ATLAS USE .....	2	Secchi Depth .....	11
		Chlorophyll <u>a</u> .....	11
SURVEILLANCE DESIGN .....	3	Phosphorus .....	12
		Nitrogen .....	12
Rationale .....	3	Reactive Silicate .....	13
Station Locations .....	3	Heterotrophic Bacteria .....	13
Sampling Procedures .....	4		
		GLOSSARY .....	14
DATA ANALYSIS .....	6		
		SYMBOLS AND ABBREVIATIONS .....	17
Regional Water Quality .....	6		
Temporal Water Quality .....	7	APPENDIX A .....	18
SIGNIFICANCE OF WATER QUALITY		APPENDIX B .....	29
VARIABLES .....	10		
		APPENDIX C .....	46
Temperature .....	10		

# INTRODUCTION

The Province of Ontario has been engaged in Lake Ontario water quality surveillance since the beginning of this century. Earliest records are contained in reports dealing with investigations of potable water supplies carried out following the signing of the 1909 Boundary Water Treaty between Great Britain and the United States of America. Beginning in 1967, an extensive nearshore monitoring network was developed by the Province leading to establishment of a comprehensive Lake Ontario water quality surveillance program. This program was set up to evaluate the effectiveness of remedial and preventive measures and to assess the suitability of the nearshore for various water uses.


In the period 1967-1975, the nearshore monitoring network consisted of stations set out along a partial one-mile grid. Representative of studies carried out during this period is information collected in 1972 which subsequently was summarized in a publication entitled "Great Lakes Water Quality Data - 1972". In order to improve the areal resolution of nearshore water quality, the original sampling grid was modified in 1976 to a transect system which, following refinement, represents the present monitoring design.

The purpose of this publication is to present the most recent data which are indicative of the trophic state of the Canadian nearshore waters of Lake Ontario. Data summarized in this Atlas span the period from 1976 to 1979 and include information on regional and temporal (annual) differences in water quality based on spring conditions. Complementary programs addressing issues other than eutrophication such as trace contaminant distribution, bacteriological water quality degradation, and problem area assessment are dealt with in separate publications. Water quality and related data collected by the Ministry of the Environment in Lake Superior, Lake Huron, Georgian Bay, Lake Erie and in the inter-connecting channels can be found in numerous other reports.

The present report should be of use to those who are interested in water quality conditions in Lake Ontario, to those who are concerned with regional nearshore developments and to those who may be undertaking studies in the area. We feel this Atlas not only provides a data base against which future changes in the nearshore can be assessed, but also serves as a useful tool for guiding future development planning along the Lake Ontario nearshore zone.

# GUIDE TO ATLAS USE

To gain an insight into the water quality conditions observed in the Lake Ontario nearshore band for any reported year, or to better comprehend the temporal trends occurring in a nearshore region of interest, read a brief description on the subject presented in sections "Regional Water Quality" and "Temporal Water Quality", and study the corresponding coloured maps and graphics. If uncertain of scientific terminology or abbreviations, look up their significance or meaning under "Significance of Water Quality Variables", "Glossary" and "Symbols and Abbreviations".



*Extensive computer facilities are used to handle a large data base generated by numerous surveillance programs.*

The following guide is intended to aid in the interpretation of information presented in the coloured maps:

Zone	water mass which is significantly different from the remaining zones
N	number of stations within each of the zones
Mean	arithmetic mean of all stations within each zone based on an average of three replicates (runs) per station (geometric mean was used in the case of heterotrophic bacteria)
SD	standard deviation of above mean
Range	Maximum and minimum station means observed within each zone.

To obtain original numerical data for areas and years of specific interest, look up corresponding station numbers presented on the transect location maps and address your parameter request to:

Ministry of the Environment  
Water Resources Branch  
Planning & Co-ordination Section  
Great Lakes Surveys Unit  
135 St. Clair Avenue West  
Toronto, Ontario, Canada  
M4V 1P5

This information is stored on the SIS data base system and is only available on microfiche.

# SURVEILLANCE DESIGN

## Rationale

Water quality in the nearshore area of Lake Ontario is exceedingly difficult to monitor because of this zone's dynamic state. In the spring a thermal bar develops, limiting exchange between the nearshore (inside the thermal bar) and offshore (beyond the thermal bar) water masses. In early summer, thermal stratification sets in, effectively isolating the epilimnetic (top) from the hypolimnetic (bottom) waters. A strongly stratified situation, however, is rare in the near-shore band since upwellings and downwellings dominate the physical processes, thus exchanging vast water masses with nearly regular frequencies. Superimposed on the macroscale dynamics, are smaller fluctuations such as seiches, which further complicate the study of this zone.

In order to appreciate the variability in water quality one must also become familiar with the seasonal changes which take place in the lake. In the winter, isothermal conditions permit complete

mixing of waters and regeneration of nutrients from sediments, with the result that in late winter, nutrient levels become elevated throughout the water column. In the spring, additional nutrients from spring runoff and rising water temperatures act to stimulate growth of diatoms. Spring chlorophyll concentrations depicted in this Atlas likely represent certain stages of the spring diatom pulse. However, since the onset, duration and intensity of the spring diatom pulse are highly variable from year to year, interpretation of chlorophyll a data based on spring conditions should be done with caution. By late spring, algal blooms largely deplete the surface (epilimnetic) waters of nutrients, while dead algal cells sink downward into the water column where they release their nutrients (through microbial action) into the hypolimnetic waters. Nutrient depletion in the epilimnion continues during the summer, when green algae predominate and further remove phosphorus and nitrogen from the waters. In the fall, the lake cools and strong winds tend to break down the remaining stratification with the result that, by early winter, the nutrient-rich hypolimnetic waters are once again mixed with the rest of the lake.

The Ministry of the Environment has developed a strategy for monitoring the nearshore areas of Lake Ontario with an aim to assessing both spatial and temporal (annual) differences while taking into account the physical processes governing this active region.

By restricting its synoptic nearshore surveillance activities to the spring period, the Ministry optimizes its monitoring effort by taking advantage of vertical homogeneity prevailing in the water column at that time, and consequently being able to restrict sampling to surface waters only. Spring sampling has the added advantage of permitting an increased resolution of nearshore regional differences in water quality since the thermal bar accentuates the effects of shore inputs by restricting their complete mixing with the open lake. Furthermore, data collected in the spring are subjected to less temporal variability than those collected in the summer months when the physical processes, such as upwellings, acting upon the nearshore region are far more disruptive to its integrity.

### Station Locations

The evolution of the nearshore transect design can be followed over the 1976-1979 period by examining station location maps. Although during this time transect locations have shifted somewhat to better resolve regional differences -and inshore-offshore gradients, the basic concept has remained unchanged.


All transects have been set up perpendicular to the shoreline with stations selectively located along depth contours of 4, 8, 12, 16, 32 and 50 metres. In 1979, there were approximately 180 stations in the nearshore area between the Niagara River mouth and Oshawa. Station locations for the 1976-1979 period are presented in Appendix B.

The offshore region of Lake Ontario is monitored by the Federal Department of the Environment.

### Sampling Procedures

Sampling commences each spring soon after the onset of the thermal bar. Surface (1.5 metres) samples are taken in triplicate runs at each of the designated stations. A table summarizing sampling periods for each survey component (run) is presented in

Appendix B. All samples are treated in compliance with standard procedures outlined in the Ministry publication entitled "A Guide to the Collection and Submission of Samples for Laboratory Analysis". In the laboratory, the samples are analyzed for a set of parameters designed to assess the trophic status of the waters. In addition, temperature, dissolved oxygen, pH and secchi depth readings are taken in the field. A summary of all parameters now available on the SIS system for the 1976-1979 period is given in Appendix B. Please note that a majority of these, but not all, have been mapped out in this Atlas.



*above*

*The Nansen bottle is often employed in the collection of water samples.*

*left*

*Certain analyses, such as dissolved oxygen, require immediate on-board determinations.*

# DATA ANALYSIS

This Atlas presents both spatial and temporal summaries of Lake Ontario nearshore water quality data.

## Regional Water Quality

Regional differences in water quality have been determined through a statistical procedure <sup>(1)</sup> which has partitioned the nearshore band into statistically different areas. A regression model was used to separate the temporal (between replicate runs) variability from spatial (between station) variability. All data, with the exception of temperature (which was plotted as isotherms), were subjected to this grouping model and subsequently presented pictorially in the accompanying maps (Appendix C).

All nearshore stations, with the exception of those located at river mouths and in small harbours, were used in the analysis. Because of their high varia-

bility, river mouth and small harbour stations were excluded from the grouping analysis. On the other hand, the two largest harbours on the lake, Toronto and Hamilton, have been included in the grouping model, by using their centrally located stations. It should be borne in mind, however, that these stations represent a conservative estimate of water quality in these two harbours and do not reflect the spatial differences existing within the harbours themselves. For further information on Toronto and Hamilton Harbours please refer to the series of Ministry reports dealing specifically with these bodies of water.

The coloured maps represent the spatial variability inherent in the nearshore areas during spring conditions. The levels of a variable in question are statistically similar within each of the designated zones while at the same time being significantly different from those in all other zones. Please note that although the mean concentrations characterizing each zone are obviously different from each other, the ranges between the zones occasionally overlap as a result of transformation procedures necessitated by the statistical analyses.

---

(1) A.H. El-Shaarawi and R.E. Kwiatkowski. A model to describe the inherent spatial and temporal variability of parameters in Lake Ontario 1974. J. Great Lakes Res., December 1977.



## Temporal Water Quality

Changes occurring within the nearshore areas of Lake Ontario over the last thirteen years (1967-1979) have been documented (Appendix A) through linear regression analyses performed on spring data collected in the following four regions: Niagara to Jordan harbour, Jordan harbour to Oakville, Oakville to Ashbridges Bay and Asbridges Bay to Ajax. These areas were chosen on the basis of regional differences observed through the statistical analyses previously discussed. The linear regression chosen to assess the temporal trends was thus based on spring arithmetic means of all stations sampled within each area with the exception of stations located at input sources. These stations were deleted from the trend analyses in order to minimize small scale localized effects and reduce variability and scatter in the data. The statistical significance of the computed regression line was tested through an analysis of variance at a 95% confidence level.

Temporal water quality trends are not only a function of the loading regime but also reflect the effect of environmental factors which govern the response of the nearshore zone. Thus, anticipated changes in

nearshore water quality, as a result of abatement measures, are not always clearly apparent due to data scatter and varying environmental conditions experienced from one year to the next. Consequently, interpretation of temporal trends should be done in conjunction with a consideration of supporting variables as well as a good understanding of factors governing the variability of the parameter in question.

For example, apparent increases in chlorophyll a may not be necessarily a function of increased phosphorus inputs but may simply reflect lower water levels or higher water temperatures prevailing at the time of sampling. Consequently, in-depth interpretations of temporal trends must take into account not only the loading regime but also such factors as precipitation, water temperatures, water levels, thermal bar advance and ultimately, the residence time of the load. A brief synopsis dealing with some of the factors governing the temporal variability of selected water quality parameters is presented in the following table to familiarize the reader with the complexity of temporal trend interpretation. Due to the non-scientific nature of this publication, only a limited number of responses to environmental factors are dealt with.



# FACTORS RESPONSIBLE FOR VARIABILITY IN SELECTED WATER QUALITY INDICATORS

RESPONSE TO ENVIRONMENTAL FACTORS				
WATER QUALITY VARIABLE	INHERENT VARIABILITY	TEMPERATURE	WATER LEVELS	PRECIPITATION
NH <sub>3</sub> , NO <sub>2</sub> , NO <sub>3</sub>	High variability, a function of the Nitrogen Cycle transformations, including N immobilization, mineralization, nitrification and denitrification.	Complex effect since temperature has a marked influence on N regeneration, turnover and assimilation processes.	Complex effect.	Complex effect.
REACTIVE SILICATE	High variability, a function of biological assimilation rates, especially in presence of diatoms.	Since this variable is biologically active, temperature has an effect on reaction rates.	During high water level years, increased shoreline erosion may contribute silicates to the water.	Complex effect.
HETEROTROPHIC BACTERIA	High variability, a function of bacterial growth patterns.	An increase in water temperature often stimulates bacterial cell division leading to higher heterotroph populations.	Complex effect.	Higher nutrient loadings associated with runoff conditions may enhance growth of heterotrophic bacteria.

# FACTORS RESPONSIBLE FOR VARIABILITY IN SELECTED WATER QUALITY INDICATORS

RESPONSE TO ENVIRONMENTAL FACTORS				
WATER QUALITY VARIABLE	INHERENT VARIABILITY	TEMPERATURE	WATER LEVELS	PRECIPITATION
CONDUCTIVITY	Moderate variability a function of inputs such as rivers, shoreline erosion etc.	Probably little effect.	During high water level years, increased shore-erosion may contribute to higher conductivity levels.	Higher conductivity often is associated with storm events.
CHLOROPHYLL <u>a</u>	Moderate to high variability, a function of algal growth patterns.	An increase in water temperature stimulates algal growth resulting in higher chlorophyll <u>a</u> levels.	Low water levels in certain instances may change the mean nearshore water depth leading to an alteration in spring warming patterns and a shift in spring algal bloom occurrence.	Higher input loadings as a result of runoff may increase levels of available nutrients and thus enhance algal growth.
SECCHI DEPTH	High variability, a function of suspended sediments, algae, etc.	Higher temperatures may stimulate algal blooms resulting in low Secchi depth readings.	During high water level years, increased shoreline erosion may contribute to high suspended solid content in the water and consequently low Secchi depth readings.	Runoff-associated loadings may contribute to higher suspended solid content and consequently lower Secchi depth readings. Also see: chlorophyll <u>a</u> .
FILTERED REACTIVE PHOSPHORUS	Moderate to high variability, partly a function of biological assimilation - excretion rates.	Since this variable is biologically active, temperature has an effect on reaction rates.	Complex effect.	Runoff-associated loadings may contribute to increased concentrations in water.

# SIGNIFICANCE OF WATER QUALITY VARIABLES

One of the chief aims of the Ministry of the Environment's nearshore surveillance program is to monitor the effectiveness of phosphorus removal programs by assessing the trophic status of the nearshore waters. This is accomplished largely through the monitoring of phosphorus and chlorophyll a levels in the nearshore region. Since phosphorus has been shown to be an important limiting factor for phytoplankton growth, cultural eutrophication, or the increase in the input rate of nutrients (notably phosphorus) from human activities, has been singled out as the prime cause of biomass increases in the Great Lakes. Cognizant of the relationship between phosphorus and eutrophication, the water management strategies encompassed in the Great Lakes Water Quality Agreement of 1978 signed by the United States of America and Canada included provisions for stringent controls on phosphorus loadings arising from human sources.

A brief description, emphasizing the importance of trophic indicators and supportive variables chosen for presentation in this Atlas, is given below. The reader is urged to supplement this discussion by referring to a booklet entitled "Water Management Goals, Policies, Objectives and Implementation Procedures of the Ministry of the Environment", published in November, 1978.

## Temperature

Water temperature is important, and at times critical, for many uses of water. It determines the solubility of gases, directly affects biological and chemical reaction rates and equilibria and governs the suitability of waters as an habitat for aquatic life. An understanding of the thermal structure prevailing during the spring period is essential for proper interpretation of water quality information presented in this Atlas. Of prime importance is the presence of the thermal bar which is defined as a 4<sup>0</sup>C transition zone between the warmer nearshore waters and the colder mid-lake waters. Since water density is greatest at 4<sup>0</sup>C, the thermal bar serves

as a density barrier which impedes mixing of nearshore water masses with the open lake. As a result, the effect of shore inputs on nearshore water quality is accentuated and plumes arising from direct discharges become well defined.

#### Conductivity (Specific Conductance)

The determination of specific conductance is a method of measuring the ionic strength of water. Ionized chemical compounds present in surface waters either naturally, or as a result of man's activities, include: calcium, magnesium, sodium, bicarbonate, carbonate, chloride, nitrate, nitrite and sulphate. Since there is a direct correlation between the total concentration of ionic species dissolved in water and specific conductance, this parameter is a sensitive, although indirect indicator of water quality changes.

#### Secchi Depth

A Secchi depth reading is a measure of water clarity. A Secchi disk is a black and white disk, 20 cms in diameter, which is lowered into the water column to a depth at which it is just visible. The Secchi depth

measurement can be used to estimate the depth to which white light penetrates the water column and can still be usable by algae for photosynthesis and growth. This usable light depth is roughly calculated to be twice the Secchi depth. Factors responsible for low Secchi depth readings are suspended and colloidal matter including sediment particles, algal cells and inorganic detritus as well as colour. Most often, low Secchi depth readings are caused by algal blooms or suspended solid inputs from erosion, landfilling, dredging, river runoff and/or industrial inputs.

#### Chlorophyll a

Chlorophyll a is a green photosynthetic pigment common to all algal groups. Because of its ubiquity in algae, chlorophyll a can be used as an indirect measure of algal densities present in the water. More specifically, chlorophyll a levels provide an appropriate measure of algal biomass which can then be used to assess the effectiveness of nutrient removal programs as well as to determine the relative trophic status of the waters. Although a universally acceptable trophic classification based on

chlorophyll a is still being developed, some authors<sup>(1)</sup> have set the following arbitrary limits as guidelines:

- eutrophic waters - chlorophyll a: >8.8 ug/L
- mesotrophic waters - chlorophyll a: 4.4 - 8.8 ug/L
- oligotrophic waters - chlorophyll a: <4.4 ug/L

Major problems associated with high algal populations include taste and odour in drinking water supplies, clogging of water intake filters, oxygen depletion, alteration of aquatic habitat, and aesthetic impairment. Only infrequent and isolated instances of these problems have taken place in Lake Ontario nearshore areas thus far.

#### Phosphorus (Total and Filtered Reactive)

Phosphorus is an essential nutrient for all living systems including algae. Total phosphorus is a measure of phosphorus content in a water sample and consists of phosphorus found in dissolved forms as well as in organic forms, including the algae themselves. To avoid nuisance concentrations of algae in lakes, it is suggested that the average total phosphorus concentrations during the ice-free

period should not generally exceed 20 ug/L. Filtered reactive phosphorus is comprised largely of ortho-phosphates, which are considered to be the forms of phosphorus available for assimilation and growth by algae. Sources of phosphorus to lakes are: untreated and treated sewage, some industrial wastes, urban stormwater, agricultural drainage, and direct precipitation.

It is widely believed that a reduction in phosphorus input into the Great Lakes will result in decreased algal populations and minimization of associated eutrophication problems. Provisions for control of phosphorus are outlined in detail in the "Great Lakes Water Quality Agreement of 1978", signed by Canada and the United States of America.

#### Nitrogen (Organic Nitrogen, Ammonia, Nitrate, Nitrite)

Nitrogen is an essential element required by all biological systems including algae. The principle forms of nitrogen occurring in aquatic ecosystems are ammonia ( $\text{NH}_3\text{-N}$ ), nitrate ( $\text{NO}_3\text{-N}$ ), nitrite ( $\text{NO}_2\text{-N}$ ) and organic nitrogen. Total Kjeldahl nitrogen includes nitrogen present in the organic form and as free ammonia.

---

(1) Dobson, H.F.H., M. Gilbertson and P.G. Sly.  
A summary and comparison of nutrients and related water quality in lakes Erie, Ontario, Huron and Superior. J. Fish Res. Board Can. 31: 731-738.

Organic nitrogen is an essential constituent of protein in all living organisms, including algae. All inorganic forms of nitrogen ( $\text{NH}_3\text{-N}$ ,  $\text{NO}_3\text{-N}$ ,  $\text{NO}_2\text{-N}$ ) are available to algae for assimilation and growth, with preferential uptake exhibited towards ammonia. The concentration of various nitrogen forms in water is the result of complex interactions, including mineralization, nitrification, and denitrification. Consequently, since the levels of inorganic nitrogen forms exhibit considerable variability, caution must be exercised in all data interpretation and especially in that pertaining to temporal trend analysis.

Principle sources of nitrogeneous compounds include municipal waste inputs (largely  $\text{NH}_3$ ), drainage from fertilized agricultural areas (mainly  $\text{NO}_2$  and  $\text{NO}_3$ ) and processed waste from chemical and steel plants ( $\text{NH}_3$ ,  $\text{NO}_2$  and  $\text{NO}_3$ ).

#### Reactive Silicate

Silicate is essential for the growth of diatoms and some species of Chrysophyceae and Xanthophyceae but

probably not for most other groups of algae. The assimilation of silicates by diatoms is directly connected with the formation of new cell walls following cell division. Spring diatom blooms tend to deplete epilimnetic waters of silica, with the result that diatoms usually do not come back into dominance until fall overturn which results in silica replenishment from hypolimnetic waters.

#### Heterotrophic Bacteria

Heterotrophic bacteria are a group of micro-organisms that require some form of organic carbon for their growth, as opposed to the rest of the bacterial community which is capable of surviving solely on inorganic substrates. Populations of heterotrophic bacteria reflect the availability of organic nutrient sources, and thus densities of these organisms in eutrophic waters are generally much higher than those found in oligotrophic waters.

# GLOSSARY

ALGAE - a group of aquatic plants which may be unicellular or multicellular and range from microscopic (most phytoplankton) to macroscopic (eg. Cladophora, a large filamentous alga which is sometimes found in mats growing or accumulating along shorelines).

AREAL - spatial

BLOOM - a sudden increase in the abundance of algal populations.

CHRYSTOPHYCEAE - a class of algae, usually brown in colour.

COLLOIDS - very small particles which tend to remain suspended and dispersed in liquids.

CONFIDENCE LEVEL - The probability (expressed as a percentage) that a tested statistical relationship is true.

DIATOMS - a class of algae which possess a cell wall consisting of two distinct halves impregnated with silica.

DOWNWELLING - see UPWELLING.

EPILIMNION - see STRATIFICATION.

EUTROPHICATION - an increase in the input rate of nutrients into a body of water with a resultant increase in biomass.

EUTROPHIC - see TROPHIC STATE.

GEOMETRIC MEAN - an antilog of the mean of logarithmically transformed data.

GRID - a network of stations forming the basis of map references.

HYPOLIMNION - see STRATIFICATION

ISOTHERM - a line joining equal values of temperatures.

LINEAR REGRESSION - relation of one variable to another expressed as a straight line

MEAN - an average defined as the sum of all observations divided by the number of observations.

MESOTROPHIC - see TROPHIC STATE.

pH - a measure of the hydrogen ion content of water on an alkalinity-acidity scale. Neutral waters have pH7, acidic less than pH7 and alkaline greater than pH7.

PLUME - a zone of influence associated with a point source discharging into a body of water.

SEICHE - a standing wave in a lake manifested by periodic oscillation of water, commonly set up by prolonged uni-directional winds.

THERMAL STATIFICATION - a partitioning of a water column into distinct thermal layers. Following spring warm-up, the lake

becomes divided into the following three regions which are strongly resistant to mixing with each other: Epilimnion - a warmer upper stratum, Thermocline - a thermal gradient zone which represents the transition between the upper warm waters and a lower cold layer, Hypolimnion - a colder lower layer.

THERMAL BAR - a narrow transition zone consisting of a nearly vertical 4° centigrade isotherm which develops in the spring and separates the warmer nearshore waters from colder open water masses.

TRANSECT - a line, perpendicular to the shoreline, along which stations have been selected.

TROPHIC STATE - a classification of water bodies based on their supply of nutrients. Eutrophic waters have a good supply of nutrients and therefore are capable of rich biological production. Mesotrophic waters are



intermediate in characteristics between eutrophic and oligotrophic waters. Oligotrophic waters are poorly supplied with nutrients and support limited biological production.

UPWELLING - a forcing of hypolimnetic waters toward the surface, brought about by the tilting of the thermocline as a result of prolonged uni-directional wind stresses. An upwelling occurring along a nearshore area nearly always takes place simultaneously with a downwelling on the opposite shore. A downwelling is a displacement of hypolimnetic waters by warmer epilimnetic waters.

XANTHOPHYCEAE - a class of algae often (yellow) green in colour.



*Well-equipped ships, such as the Guardian No.1, are used in surveillance and monitoring of the Great Lakes.*

# SYMBOLS AND ABBREVIATIONS


## QUALIFIERS





<	less than
>	greater than

## UNITS





m	meters
mg/L	milligrams per litre
ug/L	micrograms per litre
uS/cm	microsiemens per centimetre
FTU	Formazin Turbidity Units
°C	Degrees Celsius (centigrade)

## APPENDIX A









# TOTAL PHOSPHORUS TEMPORAL TRENDS IN FOUR REGIONS OF L. ONTARIO NEARSHORE







## FILTERED REACTIVE PHOSPHORUS TEMPORAL TRENDS IN FOUR REGIONS OF L. ONTARIO NEARSHORE







## ORGANIC NITROGEN

### TEMPORAL TRENDS IN FOUR REGIONS OF L. ONTARIO NEARSHORE







# NITRITE + NITRATE TEMPORAL TRENDS IN FOUR REGIONS OF L. ONTARIO NEARSHORE









## AMMONIA





### TEMPORAL TRENDS IN FOUR REGIONS OF L. ONTARIO NEARSHORE







## TOTAL KJELDAHL NITROGEN TEMPORAL TRENDS IN FOUR REGIONS OF L. ONTARIO NEARSHORE



## SECCHI DISK DEPTH TEMPORAL TRENDS IN FOUR REGIONS OF L. ONTARIO NEARSHORE




## CHLOROPHYLL a TEMPORAL TRENDS IN FOUR REGIONS OF L. ONTARIO NEARSHORE




## CONDUCTIVITY

## TEMPORAL TRENDS IN FOUR REGIONS OF L. ONTARIO NEARSHORE


## APPENDIX B




1976



1977




1978





1979


NEARSHORE AREAS OF LAKE ONTARIO SURVEYED IN THE 1976 - 1979 PERIOD





































WATER QUALITY VARIABLES SAMPLED BY THE  
MINISTRY OF THE ENVIRONMENT IN NEARSHORE AREAS OF LAKE ONTARIO  
DURING 1976-1979 SURVEY PERIODS

	YEAR					YEAR			
	1976	1977	1978	1979		1976	1977	1978	1979
Alkalinity	x	x	x	*	Nitrogen, Total Kjeldahl	x	x	x	x
Bacteria, Heterotrophic	x	*	*	*	Oxygen, Dissolved	x	x	x	x
Bacteria, Fecal Coliforms	x	*	*	*	Phenols	*	*	*	o
Bacteria, Streptococci	x	*	*	*	pH	x	x	x	x
Bacteria, Total Coliforms	x	*	*	o	Phosphorus, Total	x	x	x	x
Bacteria, Pseudomonas	x	*	*	*	Phosphorus, Reactive	x	x	x	o
Calcium	*	*	*	*	Phosphorus, Total Filtered	o	x	x	x
Carbon, Inorganic, Organic	*	*	*	*	Potassium	*	*	*	*
Chloride	x	x	*	*	Secchi Disk Depth	x	x	x	x
Chlorophyll <u>a</u>	x	o	x	x	Silicate, Reactive	x	x	x	x
Chlorophyll <u>a</u> (corrected)	x	o	x	x	Sodium	*	*	*	*
Chlorophyll <u>b</u>	x	o	x	x	Solids, Total	*	*	*	*
Conductivity	x	x	x	x	Solids, Dissolved	*	*	*	*
Fluoride	*	*	*	*	Solids, Suspended	*	*	*	*
Iron, Total	*	*	*	*	Sulphate	*	*	*	*
Magnesium	*	*	*	*	Temperature	x	x	x	x
Nitrogen, Ammonia	x	x	x	x	Turbidity	x	x	x	x
Nitrogen, Nitrite + Nitrate	x	x	x	x	Water Depths	x	x	x	x

x All Stations  
o Not Sampled  
\* Selected Stations Only

# SAMPLING DATES FOR 1976-1979 SURVEY PERIODS

1976		
RUN 1	RUN 2	RUN 3
Niagara River to Stoney Creek April 6	Stoney Creek to Niagara River April 7	Niagara River to Stoney Creek April 9
Hamilton to Etobicoke Creek April 10	Etobicoke Creek to Hamilton April 12	Hamilton to Etobicoke Creek April 13
Etobicoke Creek to Ashbridges Bay April 14	Ashbridges Bay to Etobicoke Creek April 16	Etobicoke Creek to Ashbridges Bay April 17
Scarborough to Whitby April 18	Whitby to Scarborough April 19	Scarborough to Whitby April 20
Coburg to Whitby April 24	Whitby to Coburg April 28	Coburg to Whitby April 29

1977		
RUN 1	RUN 2	RUN 3
Niagara River to Grimbsy April 10-12	Grimbsy to Niagara River April 13	Niagara River to Grimbsy April 14
Grimbsy to Credit River April 15	Credit River to Grimbsy April 16	Grimbsy to Credit River April 17
Credit River to Ashbridges Bay April 19	Ashbridges Bay to Credit River April 20	Credit River to Ashbridges Bay April 21
Scarborough to Oshawa April 23	Oshawa to Scarborough April 24	Scarborough to Oshawa April 25

SAMPLING DATES FOR 1976-1979 SURVEY PERIODS (continued)


1978		
RUN 1	RUN 2	RUN 3
Niagara River to Stoney Creek April 13-16	Stoney Creek to Niagara River April 17	Niagara River to Stoney Creek April 22
Hamilton to Credit River April 23	Credit River to Hamilton April 24	Hamilton to Credit River April 25
Lakeview to Ashbridges Bay April 26	Ashbridges Bay to Lakeview April 27	Lakeview to Ashbridges Bay April 28-30
Scarborough to Oshawa May 2	Oshawa to Scarborough May 3	Scarborough to Oshawa May 7
Bowmanville to Colborne May 9	Colborne to Bowmanville May 10	Bowmanville to Colborne May 1
Scotch Bonnet Is. to Amherst Is. May 17	Amherst Is. to Scotch Bonnet Is. May 18	Scotch Bonnet Is. to Amherst Is. May 19


1979		
RUN 1	RUN 2	RUN 3
Highland Cr. to Ontario Place April 19	Ontario Place to Highland Cr. April 28	Highland Cr. to Ontario Place April 29
Ontario Place to Clarkson April 20	Clarkson to Ontario Place April 27	Ontario Place to Clarkson April 30
Port Credit to Burlington April 21	Burlington to Port Credit April 26	Port Credit to Burlington May 1
Hamilton to Niagara River April 22-23	Niagara River to Hamilton April 24-25	Hamilton to Niagara River May 2-3








## APPENDIX C



















































































PLATE 29  
**CHLOROPHYLL *a* ( $\mu\text{g/L}$ )**  
**SPRING 1978**












































~~NOV 27 1981~~



\*96936000009564\*